

ISPM No. 15



**INTERNATIONAL STANDARDS FOR
PHYTOSANITARY MEASURES**

ISPM No. 15

***GUIDELINES FOR REGULATING WOOD PACKAGING
MATERIAL IN INTERNATIONAL TRADE***

(2002)

Produced by the Secretariat of the International Plant Protection Convention



ANNEX I

APPROVED MEASURES ASSOCIATED WITH WOOD PACKAGING MATERIAL

Heat treatment (HT)

Wood packaging material should be heated in accordance with a specific time-temperature schedule that achieves a minimum wood core temperature of 56°C for a minimum of 30 minutes³.

Kiln-drying (KD), chemical pressure impregnation (CPI), or other treatments may be considered HT treatments to the extent that these meet the HT specifications. For example, CPI may meet the HT specification through the use of steam, hot water, or dry heat.

Heat treatment is indicated by the mark HT. (see Annex II)

Methyl bromide (MB) fumigation for wood packaging material

The wood packaging material should be fumigated with methyl bromide. The treatment is indicated by the mark MB. The minimum standard for methyl bromide fumigation treatment for wood packaging material is as follows:

Temperature	Dosage rate	Minimum concentration (g/m ³) at:			
		0.5hrs.	2hrs.	4hrs.	16hrs.
21°C or above	48	36	24	17	14
16°C or above	56	42	28	20	17
11°C or above	64	48	32	22	19

The minimum temperature should not be less than 10°C and the minimum exposure time should be 16 hours.⁴

List of most significant pests targeted by HT and MB

Members of the following pest groups associated with wood packaging material are practically eliminated by HT and MB treatment in accordance with the specifications listed above:

Pest group
Insects
Anobiidae
Bostrichidae
Buprestidae
Cerambycidae
Curculionidae
Isoptera
Lyctidae (with some exceptions for HT)
Oedemeridae
Scolytidae
Siricidae
Nematodes
<i>Bursaphelenchus xylophilus</i>

³ A minimum core temperature of 56° C for a minimum of 30 min. is chosen in consideration of the wide range of pests for which this combination is documented to be lethal and a commercially feasible treatment. Although it is recognized that some pests are known to have a higher thermal tolerance, quarantine pests in this category are managed by NPPOs on a case by case basis.

⁴ Certain countries require that the minimum commodity temp should be higher.